

**IMPROVED MUSCLE-DIRECTED GENE TRANSFER BY USE OF  
RECOMBINANT AAV-1 AND AAV-6 VIRIONS**

5                                    **CROSS REFERENCE TO RELATED APPLICATIONS**

This application claims priority under 37 C.F.R. § 119(e) to Provisional Application Ser. No. 60/266,778 filed on February 6, 2001, herein incorporated by reference in its entirety.

10

**FIELD OF THE INVENTION**

11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
280  
281  
282  
283  
284  
285  
286  
287  
288  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
989  
990  
991  
992  
993  
994  
995  
996  
997  
998  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1830  
1831  
1832  
1833  
1834  
1835  
1836  
1837  
1838  
1839  
1840  
1841  
1842  
1843  
1844  
1845  
1846  
1847  
1848  
1849  
1850  
1851  
1852  
1853  
1854  
1855  
1856  
1857  
1858  
1859  
1860  
1861  
1862  
1863  
1864  
1865  
1866  
1867  
1868  
1869  
1870  
1871  
1872  
1873  
1874  
1875  
1876  
1877  
1878  
1879  
1880  
1881  
1882  
1883  
1884  
1885  
1886  
1887  
1888  
1889  
1890  
1891  
1892  
1893  
1894  
1895  
1896  
1897  
1898  
1899  
1900  
1901  
1902  
1903  
1904  
1905  
1906  
1907  
1908  
1909  
1910  
1911  
1912  
1913  
1914  
1915  
1916  
1917  
1918  
1919  
1920  
1921  
1922  
1923  
1924  
1925  
1926  
1927  
1928  
1929  
1930  
1931  
1932  
1933  
1934  
1935  
1936  
1937  
1938  
1939  
1940  
1941  
1942  
1943  
1944  
1945  
1946  
1947  
1948  
1949  
1950  
1951  
1952  
1953  
1954  
1955  
1956  
1957  
1958  
1959  
1960  
1961  
1962  
1963  
1964  
1965  
1966  
1967  
1968  
1969  
1970  
1971  
1972  
1973  
1974  
1975  
1976  
1977  
1978  
1979  
1980  
1981  
1982  
1983  
1984  
1985  
1986  
1987  
1988  
1989  
1990  
1991  
1992  
1993  
1994  
1995  
1996  
1997  
1998  
1999  
2000  
2001  
2002  
2003  
2004  
2005  
2006  
2007  
2008  
2009  
2010  
2011  
2012  
2013  
2014  
2015  
2016  
2017  
2018  
2019  
2020  
2021  
2022  
2023  
2024  
2025  
2026  
2027  
2028  
2029  
2030  
2031  
2032  
2033  
2034  
2035  
2036  
2037  
2038  
2039  
2040  
2041  
2042  
2043  
2044  
2045  
2046  
2047  
2048  
2049  
2050  
2051  
2052  
2053  
2054  
2055  
2056  
2057  
2058  
2059  
2060  
2061  
2062  
2063  
2064  
2065  
2066  
2067  
2068  
2069  
2070  
2071  
2072  
2073  
2074  
2075  
2076  
2077  
2078  
2079  
2080  
2081  
2082  
2083  
2084  
2085  
2086  
2087  
2088  
2089  
2090  
2091  
2092  
2093  
2094  
2095  
2096  
2097  
2098  
2099  
2100  
2101  
2102  
2103  
2104  
2105  
2106  
2107  
2108  
2109  
2110  
2111  
2112  
2113  
2114  
2115  
2116  
2117  
2118  
2119  
2120  
2121  
2122  
2123  
2124  
2125  
2126  
2127  
2128  
2129  
2130  
2131  
2132  
2133  
2134  
2135  
2136  
2137  
2138  
2139  
2140  
2141  
2142  
2143  
2144  
2145  
2146  
2147  
2148  
2149  
2150  
2151  
2152  
2153  
2154  
2155  
2156  
2157  
2158  
2159  
2160  
2161  
2162  
2163  
2164  
2165  
2166  
2167  
2168  
2169  
2170  
2171  
2172  
2173  
2174  
2175  
2176  
2177  
2178  
2179  
2180  
2181  
2182  
2183  
2184  
2185  
2186  
2187  
21

in order to maintain the necessary levels of the protein in the bloodstream. Even then, the concentration of protein is not maintained at physiological levels—the level of the protein is usually abnormally high immediately following the injection, and far below optimal levels prior to the injection. Additionally, injected delivery of recombinant protein often cannot deliver the protein to the target cells, tissues, or organs in the body. And, if the protein successfully reaches its target, it may be diluted to a non-therapeutic level. Furthermore, the method is inconvenient and often restricts the patient's lifestyle.

These shortcomings have fueled the desire to develop gene therapy methods for delivering sustained levels of specific proteins into patients. These methods are designed to allow clinicians to introduce deoxyribonucleic acid (DNA) coding for a nucleic acid, such as a therapeutic gene, directly into a patient (*in vivo* gene therapy) or into cells isolated from a patient or a donor (*ex vivo* gene therapy). The introduced nucleic acid then directs the patient's own cells or grafted cells to produce the desired protein product. Gene delivery, therefore, obviates the need for frequent injections. Gene therapy may also allow clinicians to select specific organs or cellular targets (e.g., muscle, blood cells, brain cells, etc.) for therapy.

DNA may be introduced into a patient's cells in several ways. There are transfection methods, including chemical methods such as calcium phosphate precipitation and liposome-mediated transfection, and physical methods such as electroporation. In general, transfection methods are not suitable for *in vivo* gene delivery. There are also methods that use recombinant viruses. Current viral-mediated gene delivery vectors include those based on retrovirus, adenovirus, herpes virus, pox virus, and adeno-associated virus (AAV). Like the retroviruses, and unlike adenovirus, AAV has the ability to integrate its genome into a host cell chromosome.

#### Adeno-Associated Virus-Mediated Gene Therapy

AAV is a parvovirus belonging to the genus Dependovirus, and has several attractive features not found in other viruses. For example, AAV can infect a wide range of host cells, including non-dividing cells. AAV can also infect cells from different species. Importantly, AAV has not been associated with any human or animal

disease, and does not appear to alter the physiological properties of the host cell upon integration. Furthermore, AAV is stable at a wide range of physical and chemical conditions, which lends itself to production, storage, and transportation requirements.

The AAV genome, a linear, single-stranded DNA molecule containing approximately 4700 nucleotides (the AAV-2 genome consists of 4681 nucleotides), generally comprises an internal non-repeating segment flanked on each end by inverted terminal repeats (ITRs). The ITRs are approximately 145 nucleotides in length (AAV-1 has ITRs of 143 nucleotides) and have multiple functions, including serving as origins of replication, and as packaging signals for the viral genome.

The internal non-repeated portion of the genome includes two large open reading frames (ORFs), known as the AAV replication (*rep*) and capsid (*cap*) regions. These ORFs encode replication and capsid gene products, respectively: replication and capsid gene products (i.e., proteins) allow for the replication, assembly, and packaging of a complete AAV virion. More specifically, a family of at least four viral proteins are expressed from the AAV *rep* region: Rep 78, Rep 68, Rep 52, and Rep 40, all of which are named for their apparent molecular weights. The AAV *cap* region encodes at least three proteins: VP1, VP2, and VP3.

In nature, AAV is a helper virus-dependent virus, i.e., it requires co-infection with a helper virus (e.g., adenovirus, herpesvirus, or vaccinia virus) in order to form functionally complete AAV virions. In the absence of co-infection with a helper virus, AAV establishes a latent state in which the viral genome inserts into a host cell chromosome or exists in an episomal form, but infectious virions are not produced. Subsequent infection by a helper virus “rescues” the integrated genome, allowing it to be replicated and packaged into viral capsids, thereby reconstituting the infectious virion. While AAV can infect cells from different species, the helper virus must be of the same species as the host cell. Thus, for example, human AAV will replicate in canine cells that have been co-infected with a canine adenovirus.

To construct infectious recombinant AAV (rAAV) containing a nucleic acid, a suitable host cell line is transfected with an AAV vector containing a nucleic acid. AAV helper functions and accessory functions are then expressed in the host cell. Once these factors come together, the HNA is replicated and packaged as though it were a

wild-type (wt) AAV genome, forming a recombinant virion. When a patient's cells are infected with the resulting rAAV, the HNA enters and is expressed in the patient's cells. Because the patient's cells lack the *rep* and *cap* genes, as well as the adenovirus accessory function genes, the rAAV are replication defective; that is, they cannot further replicate and package their genomes. Similarly, without a source of *rep* and *cap* genes, wtAAV cannot be formed in the patient's cells.

There are six known AAV serotypes, AAV-1 through AAV-6. Of those six serotypes, AAV-2 is the best characterized, having been used to successfully deliver transgenes to several cell lines, tissue types, and organs in a variety of in vitro and in vivo assays. The six serotypes of AAV can be distinguished from one another by the use of monoclonal antibodies or by employing nucleotide sequence analysis; AAV-1, AAV-2, AAV-3, and AAV-6 are 82% identical at the nucleotide level, while AAV-4 is 75 to 78% identical to the other serotypes (Russell et al. (1998) *J Virol* 72:309-319). Significant nucleotide sequence variation is noted for regions of the AAV genome that code for capsid proteins; such variable regions may be responsible for differences in serological reactivity to the capsid proteins of the various AAV serotypes.

It is known that readministration of a single AAV serotype can lead to a significant reduction in transduction efficiency. Moskalenko et al. (*J Virol* (2000) 74:176101766), for example, showed that mice with pre-existing anti-AAV-2 antibodies, when administered Factor IX in a recombinant AAV-2 virion, failed to express the Factor IX transgene, suggesting that the anti-AAV-2 antibodies blocked transduction of the rAAV-2 virion. Halbert et al. (*J Virol* (1998) 72:9795-9805) reported similar results. Others have demonstrated successful readministration of rAAV-2 virions into experimental animals, but only after immune suppression is achieved (e.g., Halbert et al., supra).

Thus, using rAAV-2 for human gene therapy is potentially problematic because anti-AAV-2 antibodies are prevalent in human populations; in fact, one study estimated that at least 80% of the general population has been infected with AAV-2 (Berns and Linden (1995) *Bioessays* 17:237-245). The identification of AAV serotypes that are not serologically cross-reactive with AAV-2 would be a significant advancement in the art. Such AAV serotypes are described herein.

## SUMMARY OF THE INVENTION

The present invention provides AAV serotypes that have the ability to efficiently transduce cell and tissue types that AAV-2 transduces poorly and/or will not be inhibited by anti-AAV-2 antibodies. In accordance with the present invention, methods and AAV vectors for use therein are provided for the efficient delivery of a heterologous nucleic acid molecule(s) (HNA) to cells or tissue of a mammal, using recombinant AAV virions. Preferably, the cells or tissue are muscle cells or muscle tissue.

More specifically, the present invention provides for the use of AAV-1 and AAV-6 serotypes (i.e., AAV virions containing AAV-1 and/or AAV-6 capsid proteins) to deliver an HNA encoding anti-sense RNA, ribozymes, or genes that express proteins, wherein expression of said anti-sense RNA, ribozymes, or genes provides for a therapeutic effect in a mammalian subject. In one embodiment, the rAAV virions containing an HNA are injected directly into a muscle. In another embodiment, the rAAV virions containing an HNA are administered into the vasculature via injection into veins, arteries, or other vascular conduits, or by using techniques such as isolated limb perfusion.

In a preferred embodiment of the invention, AAV-1-derived and AAV-6-derived virions are provided that contain a gene encoding a blood coagulation protein which, when expressed at a sufficient concentration, provides for a therapeutic effect, such effect being an improvement in the blood-clotting efficiency of a mammal suffering from a blood clotting disorder. The blood clotting disorder can be any disorder adversely affecting the organism's ability to coagulate the blood. Preferably, the blood clotting disorder is hemophilia.

In one embodiment, the gene encoding a blood coagulation protein is a Factor VIII gene. Preferably, the Factor VIII gene is the human Factor VIII gene or a derivation thereof. In another embodiment, the gene encoding a blood coagulation protein is a Factor IX gene. Preferably, the Factor IX gene is the human Factor IX (hF.IX) gene.

These and other embodiments of the instant invention will readily occur to those of ordinary skill in the art in view of the disclosure herein.

#### BRIEF DESCRIPTION OF THE DRAWINGS

5

FIG. 1 represents circulating plasma hF.IX in nanograms per milliliter (ng/mL) as measured in RAG-1 mice following intramuscular (IM) injection of  $2 \times 10^{11}$  viral vector genomes/kg (n=4).

10

FIG. 1 represents circulating plasma hF.IX in nanograms per milliliter (ng/mL) as measured in RAG-1 mice following intramuscular (IM) injection of  $2 \times 10^{11}$  viral vector genomes/kg (n=4).

## DETAILED DESCRIPTION OF THE INVENTION

The present invention embraces the use of a recombinant adeno-associated virus (rAAV) virion to deliver a “heterologous nucleic acid” (an “HNA”) to a mammalian subject. A “recombinant AAV virion” or “rAAV virion” is an infectious virus composed of an AAV protein shell (i.e., a capsid) encapsulating a “recombinant AAV (rAAV) vector,” the rAAV vector comprising the HNA and one or more AAV inverted terminal repeats (ITRs). AAV vectors can be constructed using recombinant techniques that are known in the art and include one or more HNAs flanked by functional ITRs. The ITRs of the rAAV vector need not be the wild-type nucleotide sequences, and may be altered, e.g., by the insertion, deletion, or substitution of nucleotides, so long as the sequences provide for proper function, i.e., rescue, replication, and packaging of the AAV genome.

Recombinant AAV virions may be produced using a variety of techniques known in the art, including the triple transfection method (described in detail in U.S. Patent No. 6,001,650, the entirety of which is incorporated by reference). This system involves the use of three vectors for rAAV virion production, including an AAV helper function vector, an accessory function vector, and a rAAV vector that contains the HNA. One of skill in the art will appreciate, however, that the nucleic acid sequences encoded by these vectors can be provided on two or more vectors in various combinations. As used herein, the term “vector” includes any genetic element, such as a plasmid, phage, transposon, cosmid, chromosome, artificial chromosome, virus, virion, etc., which is capable of replication when associated with the proper control elements and which can transfer gene sequences between cells. Thus, the term includes cloning and expression vehicles, as well as viral vectors.

The AAV helper function vector encodes the “AAV helper function” sequences (i.e., rep and cap), which function *in trans* for productive AAV replication and encapsidation. Preferably, the AAV helper function vector supports efficient AAV vector production without generating any detectable wild-type AAV virions (i.e., AAV virions containing functional rep and cap genes). An example of such a vector, pHLP19 is described in U.S. Patent No. 6,001,650. Another AAV helper function vector is the

pRep6cap6 vector, described in U.S. Patent No. 6,156,303, the entirety of which is herein incorporated by reference.

The accessory function vector encodes nucleotide sequences for non-AAV derived viral and/or cellular functions upon which AAV is dependent for replication (i.e., “accessory functions”). The accessory functions include those functions required for AAV replication, including, without limitation, those moieties involved in activation of AAV gene transcription, stage specific AAV mRNA splicing, AAV DNA replication, synthesis of cap expression products, and AAV capsid assembly. Viral-based accessory functions can be derived from any of the known helper viruses such as adenovirus, herpesvirus (other than herpes simplex virus type-1), and vaccinia virus. In a preferred embodiment, the accessory function plasmid pladen5 is used (details regarding pLaden5 are described in U.S. Patent No. 6,004,797, the entirety of which is hereby incorporated by reference). This plasmid provides a complete set of adenovirus accessory functions for AAV vector production, but lacks the components necessary to form replication-competent adenovirus.

The instant invention broadly contemplates the use of two specific AAV serotypes: AAV-1 and the AAV-6. The term “serotype” is used herein to describe the genotype of a virus or unicellular organism that has been defined by means of antisera binding to antigenic determinants located on the surface of the virus or unicellular organism. In the case of AAV, the antigenic determinants located on the AAV virion’s surface are the capsid proteins, so it is the capsid protein that distinguishes the AAV serotype. Capsid proteins are the product of AAV cap gene expression, the specific sequence of the cap gene being unique to the particular AAV serotype.

Recombinant AAV-6 virions are produced, in one embodiment of the invention, with an AAV helper function vector containing the rep and cap genes from the AAV-6 genome (this vector is known as pRepCap6 – see U.S. Patent No. 6,156,303, supra, for a thorough description of the pRepCap6 vector). The AAV-6 genome has been published and is available under GenBank Accession No. 9629894. One of skill in the art will also appreciate that other rep genes, e.g. rep2, can be used in combination with the cap6 gene to produce AAV “hybrid” helper function vectors, which are then capable of supporting the production of rAAV-6 virions. The term “hybrid” as used herein is an



AAV helper function vector with a rep gene from one serotype (other than AAV-6) in combination with the cap gene from the AAV-6 genome (e.g., pRep2Cap6, pRep3Cap6, etc.). The rep and cap genes can be wild-type in their sequences, or be altered by partial deletion, mutation, rearrangement, addition, gene or gene segment shuffling, etc., the primary consideration as contemplated herein being the retention of rep and cap wild-type function. See U.S. Patent No. 6,156,303, *supra*, (and Example 1 below) for methods describing the generation of rAAV-6 virions.

Similar methods can also be employed to construct AAV helper function vectors containing rep and cap1 genes. Incorporating the rep gene from the AAV-1 genome with the cap gene from the AAV-1 genome yields the AAV helper function vector pRepCap1. Incorporating a rep gene from an AAV serotype other than AAV-1 yields a hybrid AAV helper function vector still capable of supporting AAV-1 virion production since the AAV helper function vector contains the cap1 gene. Either pRepCap1 or a hybrid AAV helper function vector such as pRep2Cap1 can support rAAV-1 virion production. The AAV-1 genome has been published under the Patent Cooperation Treaty (international publication WO 0028061) and is available under GenBank Accession No. 9632547.

We have shown that the cap1 and cap6 proteins recognize the cap binding site(s) on the AAV-2 ITRs. This is thought to be because the cap1 and cap6 proteins recognize the AAV-2 ITR secondary structure, and not specific AAV-2 ITR DNA sequences. It is believed that the ITRs from the various AAV serotypes assume similar secondary structure so one of skill in the art would appreciate that AAV-1, AAV-3, AAV-4, AAV-5, or AAV-6 ITRs could be used with pRep6Cap6 or a hybrid AAV-6 helper function vector to generate AAV-6 serotype virions (or pRep1Cap1 or a hybrid AAV-1 helper function vector to generate AAV-1 serotype virions). For example, using the methods of the instant invention, pRep2Cap6 could be used in conjunction with AAV-3 ITRs to produce AAV-6 virions.

The HNA, that is, the "heterologous nucleic acid," comprises nucleic acid sequences joined together that are otherwise not found together in nature, this concept defining the term "heterologous." To illustrate the point, an example of an HNA is a gene flanked by nucleotide sequences not found in association with that gene in nature.

Another example of an HNA is a gene that itself is not found in nature (e.g., synthetic sequences having codons different from the native gene). Allelic variation or naturally occurring mutational events do not give rise to HNAs, as used herein. An HNA can comprise an anti-sense RNA molecule, a ribozyme, or a gene encoding a polypeptide.

5           The HNA is operably linked to a heterologous promoter (constitutive, cell-specific, or inducible) such that the HNA is capable of being expressed in the patient's target cells under appropriate or desirable conditions. Numerous examples of constitutive, cell-specific, and inducible promoters are known in the art, and one of skill could readily select a promoter for a specific intended use, e.g., the selection of the  
10       muscle-specific skeletal  $\alpha$ -actin promoter or the muscle-specific creatine kinase promoter/enhancer for muscle cell-specific expression, the selection of the constitutive CMV promoter for strong levels of continuous or near-continuous expression, or the selection of the inducible ecdysone promoter for induced expression. Induced expression allows the skilled artisan to control the amount of protein that is synthesized.  
15       In this manner, it is possible to vary the concentration of therapeutic product. Other examples of well known inducible promoters are: steroid promoters (e.g., estrogen and androgen promoters) and metallothionein promoters.

          The invention includes rAAV-1 or rAAV-6 virions comprising HNAs coding for one or more anti-sense RNA molecules, the rAAV virions preferably administered to  
20       one or more muscle cells or tissue of a mammal. Antisense RNA molecules suitable for use with the present invention in cancer anti-sense therapy or treatment of viral diseases have been described in the art. See, e.g., Han et al., (1991) *Proc. Natl. Acad. Sci. USA* 88:4313-4317; Uhlmann et al., (1990) *Chem. Rev.* 90:543-584; Helene et al., (1990) *Biochim. Biophys. Acta.* 1049:99-125; Agarawal et al., (1988) *Proc. Natl. Acad. Sci. USA* 85:7079-7083; and Heikkila et al., (1987) *Nature* 328:445-449. The invention also  
25       encompasses the delivery of ribozymes using the methods disclosed herein. For a discussion of suitable ribozymes, see, e.g., Cech et al., (1992) *J. Biol. Chem.* 267:17479-17482 and U.S. Pat. No. 5,225,347.

          The invention preferably encompasses rAAV-1 or rAAV-6 virions comprising  
30       HNAs coding for one or more polypeptides, the rAAV virions preferably administered to one or more muscle cells or tissue of a mammal. Thus, the invention embraces the

delivery of HNAs encoding one or more peptides, polypeptides, or proteins, which are useful for the treatment of disease states in a mammalian subject. Such DNA and associated disease states include, but are not limited to: DNA encoding glucose-6-phosphatase, associated with glycogen storage deficiency type 1A; DNA encoding phosphoenolpyruvate-carboxykinase, associated with Pepck deficiency; DNA encoding galactose-1 phosphate uridyl transferase, associated with galactosemia; DNA encoding phenylalanine hydroxylase, associated with phenylketonuria; DNA encoding branched chain alpha-ketoacid dehydrogenase, associated with Maple syrup urine disease; DNA encoding fumarylacetoacetate hydrolase, associated with tyrosinemia type 1; DNA encoding methylmalonyl-CoA mutase, associated with methylmalonic acidemia; DNA encoding medium chain acyl CoA dehydrogenase, associated with medium chain acetyl CoA deficiency; DNA encoding ornithine transcarbamylase, associated with ornithine transcarbamylase deficiency; DNA encoding argininosuccinic acid synthetase, associated with citrullinemia; DNA encoding low density lipoprotein receptor protein, associated with familial hypercholesterolemia; DNA encoding UDP-glucouronosyltransferase, associated with Crigler-Najjar disease; DNA encoding adenosine deaminase, associated with severe combined immunodeficiency disease; DNA encoding hypoxanthine guanine phosphoribosyl transferase, associated with Gout and Lesch-Nyan syndrome; DNA encoding biotinidase, associated with biotinidase deficiency; DNA encoding beta-glucocerebrosidase, associated with Gaucher disease; DNA encoding beta-glucuronidase, associated with Sly syndrome; DNA encoding peroxisome membrane protein 70 kDa, associated with Zellweger syndrome; DNA encoding porphobilinogen deaminase, associated with acute intermittent porphyria; DNA encoding alpha-1 antitrypsin for treatment of alpha-1 antitrypsin deficiency (emphysema); DNA encoding erythropoietin for treatment of anemia due to thalassemia or to renal failure; DNA encoding vascular endothelial growth factor, DNA encoding angiopoietin-1, and DNA encoding fibroblast growth factor for the treatment of ischemic diseases; DNA encoding thrombomodulin and tissue factor pathway inhibitor for the treatment of occluded blood vessels as seen in, for example, atherosclerosis, thrombosis, or embolisms; DNA encoding aromatic amino acid decarboxylase (AADC), and DNA encoding tyrosine hydroxylase (TH) for the treatment of Parkinson's disease;

DNA encoding the beta adrenergic receptor, DNA encoding anti-sense to, or DNA encoding a mutant form of, phospholamban, DNA encoding the sarco(endo)plasmic reticulum adenosine triphosphatase-2 (SERCA2), and DNA encoding the cardiac adenylyl cyclase for the treatment of congestive heart failure; DNA encoding a tumor suppressor gene such as p53 for the treatment of various cancers; DNA encoding a cytokine such as one of the various interleukins for the treatment of inflammatory and immune disorders and cancers; DNA encoding dystrophin or minidystrophin and DNA encoding utrophin or miniutrophin for the treatment of muscular dystrophies; and, DNA encoding insulin for the treatment of diabetes.

The invention also includes rAAV-1 and rAAV-6 virions comprising a gene or genes coding for blood coagulation proteins, which proteins may be delivered, using the methods of the present invention, to the cells of a mammal having hemophilia for the treatment of hemophilia. Thus, the invention includes: delivery of the Factor IX gene to a mammal for treatment of hemophilia B, delivery of the Factor VIII gene to a mammal for treatment of hemophilia A, delivery of the Factor VII gene for treatment of Factor VII deficiency, delivery of the Factor X gene for treatment of Factor X deficiency, delivery of the Factor XI gene for treatment of Factor XI deficiency, delivery of the Factor XIII gene for treatment of Factor XIII deficiency, and, delivery of the Protein C gene for treatment of Protein C deficiency. Delivery of each of the above-recited genes to the cells of a mammal is accomplished by first generating a rAAV virion comprising the gene and then administering the rAAV virion to the mammal. Thus, the invention includes rAAV virions comprising genes encoding any one of Factor IX, Factor VIII, Factor X, Factor VII, Factor XI, Factor XIII or Protein C.

Delivery of rAAV-1 or rAAV-6 virions containing one or more HNAs to a mammalian subject may be by intramuscular injection or by administration into the bloodstream of the mammalian subject. Administration into the bloodstream may be by injection into a vein, an artery, or any other vascular conduit such as a venule, an arteriole, or capillary. Additionally, a skilled artisan can administer rAAV-1 or rAAV-6 virions into the bloodstream by way of isolated limb perfusion, a technique well known in the surgical arts, the method essentially enabling the artisan to isolate a limb from the systemic circulation prior to administration of the rAAV virions. A variant of the

isolated limb perfusion technique, described in U.S. Patent No. 6,177,403 and herein incorporated by reference, can also be employed by the skilled artisan to administer rAAV-1 or rAAV-6 virions into the vasculature of an isolated limb to potentially enhance transduction into muscle cells or tissue.

5           The dose of rAAV virions required to achieve a particular “therapeutic effect,” e.g., the units of dose in vector genomes/per kilogram of body weight (vg/kg), will vary based on several factors including, but not limited to: the route of rAAV virion administration, the level of gene (or anti-sense RNA or ribozyme) expression required to achieve a therapeutic effect, the specific disease or disorder being treated, a host  
10 immune response to the rAAV virion, a host immune response to the gene (or anti-sense RNA or ribozyme) expression product, and the stability of the gene (or anti-sense RNA or ribozyme) product. One of skill in the art can readily determine a rAAV virion dose range to treat a patient having a particular disease or disorder based on the aforementioned factors, as well as other factors that are well known in the art.

15           Generally speaking, by “therapeutic effect” is meant a level of expression of one or more HNAs sufficient to alter a component of a disease (or disorder) toward a desired outcome or clinical endpoint, such that a patient’s disease or disorder shows clinical improvement, often reflected by the amelioration of a clinical sign or symptom relating to the disease or disorder. Using hemophilia as a specific disease example, a  
20 “therapeutic effect” for hemophilia is defined herein as an increase in the blood-clotting efficiency of a mammal afflicted with hemophilia, efficiency being determined, for example, by well known endpoints or techniques such as employing assays to measure whole blood clotting time or activated prothromboplastin time. Reductions in either whole blood clotting time or activated prothromboplastin time are indications of an  
25 increase in blood-clotting efficiency. In severe cases of hemophilia, hemophiliacs having less than 1% of normal levels of Factor VIII or Factor IX have a whole blood clotting time of greater than 60 minutes as compared to approximately 10 minutes for non-hemophiliacs. Expression of 1% or greater of Factor VIII or Factor IX has been shown to reduce whole blood clotting time in animal models of hemophilia, so  
30 achieving a circulating Factor VIII or Factor IX plasma concentration of greater than

1% will likely achieve the desired therapeutic effect of an increase in blood-clotting efficiency.

Rather than focusing exclusively on treating a disease, it is often desirable to deliver an HNA to a host cell in order to elucidate its physiological or biochemical function(s). The HNA can be either an endogenous gene or heterologous. Using either an ex vivo or in vivo approach, the skilled artisan can administer rAAV-1 and/or rAAV-6 virions containing one or more HNAs of unknown function to an experimental animal, express the HNA(s), and observe any subsequent functional changes. Such changes can include: protein-protein interactions, alterations in biochemical pathways, alterations in the physiological functioning of cells, tissues, organs, or organ systems, and/or the stimulation or silencing of gene expression.

Alternatively, the skilled artisan can over-express a gene of known function and examine its effects. Such genes can be either endogenous to the experimental animal or heterologous in nature (i.e., a transgene).

By using the methods of the present invention, the skilled artisan can also abolish or significantly reduce gene expression, thereby employing another means of determining gene function. One method of accomplishing this is by way of administering antisense RNA-containing rAAV virions to an experimental animal, expressing the antisense RNA molecule so that the targeted endogenous gene is “knocked out,” and then observing any subsequent physiological or biochemical changes.

The methods of the present invention are compatible with other well-known technologies such as transgenic mice and knockout mice and can be used to complement these technologies. One skilled in the art can readily determine combinations of known technologies with the methods of the present invention to obtain useful information on gene function.

Once delivered, in many instances it is not enough to simply express the HNA; instead, it is often desirable to vary the levels of HNA expression. Varying HNA expression levels, which varies the dose of the HNA expression product, is frequently useful in acquiring and/or refining functional information on the HNA. This can be accomplished, for example by incorporating a heterologous inducible promoter into the

5 rAAV virion containing the HNA so that the HNA will be expressed only when the promoter is induced. Some inducible promoters can also provide the capability for refining HNA expression levels; that is, varying the concentration of inducer will fine-tune the concentration of HNA expression product. This is sometimes more useful than having an “on-off” system (i.e., any amount of inducer will provide the same level of HNA expression product, an “all or none” response). Numerous examples of inducible promoters are known in the art including the ecdysone promoter, steroid promoters (e.g., estrogen and androgen promoters) and metallothionein promoters.

10 The methods of the present invention can be used to facilitate pharmaco- or toxico-kinetic studies. For example, because AAV is known to transduce hepatocytes with high efficiency, human metabolic enzymes (e.g., various oxidases and reductases such as the cytochrome p450 isozymes, various epoxide hydrolases, various dehydrogenases such as alcohol and aldehyde dehydrogenases, various peptidases, etc. – metabolic enzymes that are expressed and function in hepatocytes) can be delivered to the liver of mice by way of rAAV-1 and or rAAV-6 virions, expressed, and then various drugs and/or toxicants can be administered to the transduced mice in order to screen for any metabolites of interest.

20 The presented methods resulted in an unexpected transduction efficiency of rAAV-6 in the skeletal muscle of mice, with transduction efficiency measured by circulating plasma levels of hF.IX (the hF.IX gene delivered to the skeletal muscle of mice by rAAV-6 virions). As shown in FIG. 1, after three weeks post-injection of rAAV-6-hF.IX, serum levels of hF.IX were approximately 32-fold greater than serum levels of hF.IX in mice injected with rAAV2-hF.IX virions. After seven weeks following injection, hF.IX delivered by rAAV-6 remained at higher concentrations than rAAV-2-delivered hF.IX. For rAAV-1-hF.IX, circulating hF.IX levels were approximately 18-fold higher than circulating F.IX levels obtained from rAAV-2-hF.IX mice. After eleven weeks post-injection, the difference between rAAV-1-hF.IX and rAAV-2-hF.IX increased to 50-fold.

30 The following examples are presented in order to more fully illustrate the preferred embodiments of the invention. They should in no way be construed, however, as limiting the broad scope of the invention.

EXAMPLE 1  
RECOMBINANT AAV FACTOR IX VIRION PREPARATION

5           Recombinant AAV virions containing the human Factor IX (hF.IX) gene – the complete cDNA sequence for hF.IX available under GenBank Accession No. 182612 – were prepared using a triple-transfection procedure described in U.S. Patent No. 6,001,650, supra.

Vector Construction

AAV pRepCap6 Helper Function Vector

10           The pRepCap6 AAV helper function vector was constructed using standard molecular biological techniques. Using an infectious AAV-6 virion (the published wild-type sequence available under GenBank Accession No. 9629894), two Bgl II restriction sites were engineered into the AAV-6 genome, one just upstream of the p5  
15           promoter and one downstream of the polyadenylation site, creating the pAAV6Bgl plasmid. The Bgl II fragment containing the rep6 and cap6 genes was excised from pAAV6Bgl and inserted into a pBLUE-SCRIPT (Stratagene, La Jolla, CA) backbone to create the AAV-6 helper vector pRepCap6.

pLadeno5 Accessory Function Vector

20           The accessory function vector pLadeno5 was constructed as follows: DNA fragments encoding the E2a, E4, and VA RNA regions isolated from purified adenovirus serotype-2 DNA (obtained from Gibco/BRL) were ligated into a plasmid called pAmpscript. The pAmpscript plasmid was assembled as follows: oligonucleotide-directed mutagenesis was used to eliminate a 623-bp region including  
25           the polylinker and alpha complementation expression cassette from pBSII s/k+ (obtained from Stratagene), and replaced with an EcoRV site. The sequence of the mutagenic oligo used on the oligonucleotide-directed mutagenesis was 5'-CCGCTACAGGGCGCGATATCAGCTCACTCAA-3'. A polylinker (containing the following restriction sites: Bam HI; KpnI; SrfI; XbaI; ClaI; Bst1107I; SalI; PmeI; and  
30           NdeI) was synthesized and inserted into the EcoRV site created above such that the BamHI side of the linker was proximal to the fl origin in the modified plasmid to



provide the pAmpscript plasmid. The sequence of the polylinker was 5'-GGATCCGGTACCGCCCGGGCTCTAGAATCGATGTATACGTCGACGTTTAAACCATATG-3'.

DNA fragments comprising the adenovirus serotype-2 E2a and VA RNA sequences were cloned directly into pAmpscript. In particular, a 5962-bp SrfI-KpnI(partial) fragment containing the E2a region was cloned between the SrfI and KpnI sites of pAmpscript. The 5962-bp fragment comprises base pairs 21,606-27,568 of the adenovirus serotype-2 genome. The complete sequence of the adenovirus serotype-2 genome is accessible under GenBank No. 9626158.

The DNA comprising the adenovirus serotype-2 E4 sequences had to be modified before it could be inserted into the pAmpscript polylinker. Specifically, PCR mutagenesis was used to replace the E4 proximal, adenoviral terminal repeat with a SrfI site. The location of this SrfI site is equivalent to base pairs 35,836-35,844 of the adenovirus serotype-2 genome. The sequences of the oligonucleotides used in the mutagenesis were: 5'-AGAGGCCCGGGCGTTTTAGGGCGGAGTAACTTGC-3' and 5'-ACATACCCGCAGGCGTAGAGAC-3'. A 3,192 bp E4 fragment, produced by cleaving the above-described modified E4 gene with SrfI and SpeI, was ligated between the SrfI and XbaI sites of pAmpscript which already contained the E2a and VA RNA sequences to result in the pLaden5 plasmid. The 3,192-bp fragment is equivalent to base pairs 32,644-35,836 of the adenovirus serotype-2 genome.

#### rAAV-2 hF.IX vector

The rAAV-2 hF.IX vector is an 11,442-bp plasmid containing the cytomegalovirus (CMV) immediate early promoter, exon 1 of hF.IX, a 1.4-kb fragment of hF.IX intron 1, exons 2-8 of h.FIX, 227 bp of h.FIX 3' UTR, and the SV40 late polyadenylation sequence between the two AAV-2 inverted terminal repeats (U.S. Patent No. 6,093,392, herein incorporated by reference). The 1.4-kb fragment of hF.IX intron 1 consists of the 5' end of intron 1 up to nucleotide 1098 and the sequence from nucleotide 5882 extending to the junction with exon 2. The CMV immediate early promoter and the SV40 late polyadenylation signal sequences can be obtained from the published sequence of pCMV-Script®, which is available from the Stratagene catalog, Stratagene, La Jolla, CA, and from their website, [www.stratagene.com](http://www.stratagene.com).

### Triple Transfection Procedure

Specifically, for rAAV-6 virion production, the AAV helper function rep<sup>6</sup>cap<sup>6</sup> vector (described in U.S. Patent No. 6,156,303, supra), the accessory function vector pLadeno5 (described in U.S. Patent No. 6,004,797, supra), and the rAAV2-hF.IX vector (U.S. Patent No. 6,093,392, supra) were used. Briefly, human embryonic kidney cells type 293 (293 cells – available from the American Type Culture Collection, catalog number CRL-1573) were seeded in 10 cm tissue culture-treated sterile dishes at a density of  $3 \times 10^6$  cells per dish in 10 mL of cell culture medium consisting of Dulbecco's modified Eagle's medium supplemented with 10% fetal calf serum and incubated in a humidified environment at 37° C in 5% CO<sub>2</sub>. After overnight incubation, 293 cells were approximately eighty-percent confluent. The 293 cells were then transfected with DNA by the calcium phosphate precipitate method, a transfection method well known in the art. Briefly, 10 µg of each vector (pRepCap6, pLadeno5, and rAAV2-hF.IX) were added to a 3-mL sterile, polystyrene snap cap tube using sterile pipette tips. 1.0 mL of 300 mM CaCl<sub>2</sub> (JRH grade) was added to each tube and mixed by pipetting up and down. An equal volume of 2X HBS (274 mM NaCl, 10 mM KCl, 42 mM HEPES, 1.4 mM Na<sub>2</sub>PO<sub>4</sub>, 12 mM dextrose, pH 7.05, JRH grade) was added with a 2-mL pipette, and the solution was pipetted up and down three times. The DNA mixture was immediately added to the 293 cells, one drop at a time, evenly throughout the dish. The cells were then incubated in a humidified environment at 37° C in 5% CO<sub>2</sub> for six hours. A granular precipitate was visible in the transfected cell cultures. After six hours, the DNA mixture was removed from the cells, which were then provided with fresh cell culture medium and incubated for an additional 72 hours.

After 72 hours, the cells were lysed and then treated with nuclease to reduce residual cellular and plasmid DNA. After precipitation, rAAV virions were purified by two cycles of isopycnic centrifugation; fractions containing rAAV virions were pooled, dialysed, and concentrated. The concentrated virions were formulated, sterile filtered (0.22 µM) and aseptically filled into glass vials. Vector genomes were quantified by the “Real Time Quantitative Polymerase Chain Reaction” method (Real Time Quantitative PCR. Heid C.A., Stevens J., Livak K.J., and Williams P.M. 1996. Genome Research 6:986-994. Cold Spring Harbor Laboratory Press).

Recombinant AAV1-hF.IX virions were produced in an analogous manner to rAAV6-hF.IX virions, with a pRep1Cap1 AAV helper function vector used in place of the pRepCap6 AAV helper function vector.

Recombinant AAV-2 virions were produced with the pHLP19 helper function vector (described in U.S. Patent No. 6,001,650, supra), the pLadeno5 plasmid, and the rAAV2-hF.IX expression plasmid.

## EXAMPLE 2

### HEMOPHILIA B TREATMENT IN RAG-1 MICE WITH rAAV1-HF.IX, rAAV2-HF.IX, AND rAAV-6-HF.IX

RAG-1 female immunodeficient mice (homozygous for a mutation in the recombinase activating gene 1, and functionally equivalent to severe combined immunodeficiency mice because these mice do not produce mature B or T cells) 4-6 weeks old (obtained from Jackson Laboratories, Bar Harbor, ME) were injected with rAAV-6 virions (prepared as described in Example 1). Mice were anesthetized with an intraperitoneal injection of ketamine (70 mg/kg) and xylazine (10 mg/kg), and a 1 cm longitudinal incision was made in the lower extremity. Recombinant AAV6-hF.IX ( $2 \times 10^{11}$  viral vector genomes/kg in HEPES-Buffered-Saline, pH 7.8) virions were injected into the tibialis anterior (25  $\mu$ L) and the quadriceps muscle (50  $\mu$ L) of each leg using a Hamilton syringe. Incisions were closed with 4-0 Vicryl suture. Blood samples were collected at seven-day intervals from the retro-orbital plexus in microhematocrit capillary tubes and plasma assayed for hF.IX by ELISA. Human F.IX antigen in mouse plasma was assessed by ELISA as described by Walter et al. (*Proc Natl Acad Sci USA* (1996) 3:3056-3061). The ELISA did not cross-react with mouse F.IX. All samples were assessed in duplicate. Protein extracts obtained from injected mouse muscle were prepared by maceration of muscle in PBS containing leupeptin (0.5 mg/mL) followed by sonication. Cell debris was removed by microcentrifugation, and 1:10 dilutions of the protein extracts were assayed for hF.IX in the ELISA. The circulating plasma concentrations of hF.IX, as measured by ELISA after three weeks post-IM injection, were 185 ng/mL for rAAV-6 hF.IX gene delivery, 110 ng/mL for rAAV1-hF.IX gene

delivery, and 6 ng/mL for rAAV-2 hF.IX gene delivery. After seven weeks post-injection, hF.IX plasma concentrations increased to approximately 190 ng/mL for rAAV6- hF.IX gene delivery, 200 ng/mL for rAAV1-hF.IX gene delivery, and 20 ng/mL for rAAV2-hF.IX gene delivery (see FIG. 1). After eleven weeks post-injection, hF.IX plasma concentrations increased to approximately 300 ng/mL for rAAV1-hF.IX gene delivery, but decreased to approximately 10 ng/mL for rAAV2-hF.IX gene delivery.

### EXAMPLE 3

#### HEMOPHILIA B TREATMENT IN DOGS WITH AAV1-cF.IX

A colony of dogs having severe *hemophilia* B comprising males that are hemizygous and females that are homozygous for a point mutation in the catalytic domain of the canine factor IX (cF.IX) gene, was used to test the efficacy of cF.IX delivered by rAAV-1 virions (rAAV1-cF.IX). The severe hemophilic dogs lack plasma cF.IX, which results in an increase in whole blood clotting time (WBCT) to >60 minutes (normal dogs have a WBCT between 6-8 minutes), and an increase in activated partial thromboplastin time (aPTT) to 50-80 seconds (normal dogs have an aPTT between 13-18 seconds). These dogs experience recurrent spontaneous hemorrhages. Typically, significant bleeding episodes are successfully managed by the single intravenous infusion of 10 mL/kg of normal canine plasma; occasionally, repeat infusions are required to control bleeding.

Under general anesthesia, hemophilia B dogs were injected intramuscularly with rAAV1-cF.IX virions at a dose of  $1 \times 10^{12}$  vg/kg. The animals were not given normal canine plasma during the procedure.

Whole blood clotting time was assessed for cF.IX in plasma. Activated partial thromboplastin time was measured. A coagulation inhibitor screen was also performed. Plasma obtained from a treated hemophilic dog and from a normal dog was mixed in equal volumes and was incubated for 2 hours at 37° C. The inhibitor screen was scored as positive if the aPTT clotting time was 3 seconds longer than that of the controls (normal dog plasma incubated with imidazole buffer and pre-treatment hemophilic dog

plasma incubated with normal dog plasma). Neutralizing antibody titer against AAV vector was assessed.

In the hemophilia B dogs injected with AAV1-cF.IX, WBCT was shortened from > 60 min to 13 min (normal: 12-15 min).

5

#### EXAMPLE 4

##### HEMOPHILIA B TREATMENT IN HUMANS WITH AAV6-HF.IX

10 On Day 0 of the protocol patients are infused with hF.IX concentrate to bring factor levels up to ~100%, and, under ultrasound guidance, rAAV6-h.FIX virions are injected directly into 10-12 sites in the *vastus lateralis* of either or both anterior thighs. Injectate volume at each site is 250-500  $\mu$ L, and sites are at least 2 cm apart. Local anesthesia to the skin is provided by ethyl chloride or eutectic mixture of local  
15 anesthetics. To facilitate subsequent muscle biopsy, the skin overlying several injection sites is tattooed and the injection coordinates recorded by ultrasound. Patients are observed in the hospital for 24 h after injection; routine isolation precautions will be observed during this period to minimize any risk of horizontal transmission of virions. Patients are discharged and seen daily in outpatient clinic daily for three days after  
20 discharge, then weekly at the home hemophilia center for the next eight weeks, then twice monthly up to five months, then monthly for the remainder of the year, then annually in follow-up. Circulating plasma levels of hF.IX are quantified using ELISA as described in Example 2.